

SURGERY GRAND ROUNDS
Self Assessment Answer Sheet
Department of Surgery, University of Louisville
Auditorium, Lower Level, Ambulatory Care Building
7:00 – 8:00 a.m.

Friday, December 9, 2011

Charles B. Ross, M.D., F.A.C.S.

Associate Professor of Surgery and Chief

Division of Vascular Surgery and Endovascular Therapeutics, University of Louisville

1. **Acute thrombosis of the inferior vena cava may be manifested by symptoms including lower abdominal and/or back pain, syncope, and hypotension with or without obvious lower extremity swelling. *True or False***

ANSWER: True

2. **Absolute indications for placement of inferior vena caval filters include recurrent pulmonary embolism despite adequate anticoagulation therapy and pulmonary embolism in patients in whom anticoagulation therapy is contraindicated. *True or False***

ANSWER: True

3. **Complications of inferior vena caval filters include caval thrombosis, migration, perforation, and strut fracture with embolization? *True or False***

ANSWER: True

4. **Placement of retrievable filters and their subsequent removal once the threat of pulmonary embolism has passed is recommended by the FDA? *True or False***

ANSWER: True

5. **Endovascular techniques including catheter-directed thrombolysis and pharmacomechanical thrombectomy are indicated only for the treatment of phlegmasia cerulea dolens with impending venous gangrene? *True or False***

ANSWER: False

6. **Stenting of the iliac veins is contraindicated in females of childbearing age? *True or False***

ANSWER: False

1. Acute thrombosis of the inferior vena cava may be manifested by symptoms including lower abdominal and/or back pain, syncope, and hypotension with or without obvious lower extremity swelling. True or False
2. Absolute indications for placement of inferior vena caval filters include recurrent pulmonary embolism despite adequate anticoagulation therapy and pulmonary embolism in patients in whom anticoagulation therapy is contraindicated. True or False
3. Complications of inferior vena caval filters include caval thrombosis, migration, perforation, and [strut fracture with embolization](#)? True or False
4. Placement of retrievable filters and their subsequent removal once the threat of pulmonary embolism has passed is recommended by the FDA? True or False
5. Endovascular techniques including catheter-directed thrombolysis and pharmacomechanical thrombectomy are indicated only for the treatment of phlegmasia cerulea dolens with impending venous gangrene? True or False
6. Stenting of the iliac veins is contraindicated in females of childbearing age? True or False